

BMI1

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW: 37kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human BMI1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

fl

Synonym:

BMI1 ; Polycomb complex protein BMI 1; RING finger protein 51; RNF51; PCGF4

Catalog #: A0211

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 648

Isotype: IgG

Swiss Prot: P35226

Purity: Affinity purification

For research use only.

Background:

The polycomb group (PcG) of proteins contributes to the maintenance of cell identity, stem cell self-renewal, cell cycle regulation, and oncogenesis by maintaining the silenced state of genes that promote cell lineage specification, cell death, and cell-cycle arrest (1-4). PcG proteins exist in two complexes that cooperate to maintain long-term gene silencing through epigenetic chromatin modifications. The first complex, EED-EZH2, is recruited to genes by DNA-binding transcription factors and methylates histone H3 on Lys27. This histone methyl-transferase activity requires the Ezh2, Eed, and Suz12 subunits of the complex (5). Histone H3 methylation at Lys27 facilitates the recruitment of the second complex, PRC1, which ubiquitinylates histone H2A on Lys119 (6). Bmi1 is a component of the PRC1 complex, which together with Ring1 strongly enhances the E3 ubiquitin ligase activity of the Ring2 catalytic subunit (7). Bmi1 plays an important role in the regulation of cell proliferation and senescence through repression of the p16 INK4A and p19 ARF genes and is required for maintenance of adult hematopoietic and neural stem cells (3,4,8-10).

To place an order, please [Click HERE](#).