

CDH2

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:1000 IHC 1:50 - 1:100

Calculated MW: 100kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant Protein of human CDH2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Concentration:

1 mg/ml

Synonym:

CDH2; CD325 ; CDHN ; CDw325; NCAD; Cadherin-2; CDw325 ; Neural cadherin ;

Catalog #: A0432

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 1000

Isotype: IgG

Swiss Prot: P19022

Purity: Affinity purification

For research use only.

Background:

, R-, B- and E-cadherins, as well as about ten other members that are found in adherens junctions, a cellular structure near the apical surface of polarized epithelial cells. The cytoplasmic domain of classical cadherins interacts with β -catenin, γ -catenin (also called plakoglobin), and p120 catenin. β -catenin and γ -catenin associate with α -catenin, which links the cadherin-catenin complex to the actin cytoskeleton (1,2). While β - and γ -catenin play structural roles in the junctional complex, p120 regulates cadherin adhesive activity and trafficking (1-4). E-cadherin is considered an active suppressor of invasion and growth of many epithelial cancers (1-3). Recent studies indicate that cancer cells have up-regulated N-cadherin in addition to loss of E-cadherin. This change in cadherin expression is called the "cadherin switch". N-cadherin cooperates with the FGF receptor, leading to overexpression of MMP-9 and cellular invasion (3). In endothelial cells, VE-cadherin signaling, expression, and localization correlate with vascular permeability and tumor angiogenesis (5,6). Expression of P-cadherin, which is normally present in epithelial cells, is also altered in ovarian and other human cancers (7,8).

To place an order, please [Click HERE](#).