

NCF4

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:200 - 1:1000 IHC 1:20 - 1:200

Calculated MW: 39kDa

Observed MW: Refer to figures

Immunogen:

A synthetic Peptide of human NCF4

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Catalog #: A0610

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 4689

Isotype: IgG

Swiss Prot: Q15080

Purity: Affinity purification

For research use only.

Background:

The protein encoded by this gene is a cytosolic regulatory component of the superoxide-producing phagocyte NADPH-oxidase, a multicomponent enzyme system important for host defense. This protein is preferentially expressed in cells of myeloid lineage. It interacts primarily with neutrophil cytosolic factor 2 (NCF2/p67-phox) to form a complex with neutrophil cytosolic factor 1 (NCF1/p47-phox), which further interacts with the small G protein RAC1 and translocates to the membrane upon cell stimulation. This complex then activates flavocytochrome b, the membrane-integrated catalytic core of the enzyme system. The PX domain of this protein can bind phospholipid products of the PI(3) kinase, which suggests its role in PI(3) kinase-mediated signaling events. The phosphorylation of this protein was found to negatively regulate the enzyme activity. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008]

To place an order, please [Click HERE](#).