

LCN2

Reactivity: Human

Tested applications: WB IF

Recommended Dilution: WB 1:500 - 1:2000 IF 1:10 - 1:100

Calculated MW: 23kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human LCN2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

LCN2; 24p3; MSFI; NGAL; Lipocalin-2;

Catalog #: A11130

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 3934

Isotype: IgG

Swiss Prot: P80188

Purity: Affinity purification

For research use only.

Background:

Lipocalin-2, a member of the lipocalin family of proteins, was originally identified as a gelatinase-associated component of neutrophil secretory granules (1). Lipocalin-2 is involved in innate immunity, iron homeostasis, and apoptosis. Lipocalin-2 limits bacterial growth by binding to bacterial siderophores and sequestering iron (2-4). In mammalian cells, iron-loaded lipocalin-2 binds to its receptor, 24p3R, and is internalized, thereby releasing iron and increasing the intracellular iron concentration (5). Conversely, iron-free lipocalin-2 promotes apoptosis (5). Lipocalin-2 is also expressed in adipose tissue and promotes insulin resistance in cultured mouse adipocytes (6).

To place an order, please [Click HERE](#).