

MMP2

Reactivity: Human Mouse Rat

Tested applications: WB

Recommended Dilution: WB 1:500 - 1:2000

Calculated MW: 74kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human MMP2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CLG4; MONA; CLG4A; TBE-1; MMP-II;

Catalog #: A11144

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 4313

Isotype: IgG

Swiss Prot: P08253

Purity: Affinity purification

For research use only.

Background:

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades type IV collagen, the major structural component of basement membranes. The enzyme plays a role in endometrial menstrual breakdown, regulation of vascularization and the inflammatory response. Mutations in this gene have been associated with Winchester syndrome and Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Two transcript variants encoding different isoforms have been found for this gene.

To place an order, please [Click HERE](#).