

NOTCH1

Reactivity: Human Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:100 - 1:200

Calculated MW: 273kDa

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human NOTCH1

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

hN1; AOS5; TAN1; AOVD1;

Catalog #: A11171

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 4851

Isotype: IgG

Swiss Prot: P46531

Purity: Affinity purification

For research use only.

Background:

This gene encodes a member of the Notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. The Notch signaling network is an evolutionarily conserved intercellular signaling pathway which regulates interactions between physically adjacent cells. In *Drosophila*, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signaling pathway that plays a key role in development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remain to be determined. This protein is cleaved in the trans-Golgi network, and presented on the cell surface as a heterodimer. This protein functions as a receptor for membrane bound ligands, and may play multiple roles during development.

To place an order, please [Click HERE](#).