

PTEN

Reactivity: Human

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:100

Calculated MW: 47kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human PTEN

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

BZS; DEC; GLM2; MHAM; TEP1; MMAC1; PTEN1; 10q23del;

Catalog #: A11189

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 5728

Isotype: IgG

Swiss Prot: P60484

Purity: Affinity purification

For research use only.

Background:

PTEN (phosphatase and tensin homologue deleted on chromosome ten), also referred to as MMAC (mutated in multiple advanced cancers) phosphatase, is a tumor suppressor implicated in a wide variety of human cancers (1). PTEN encodes a 403 amino acid polypeptide originally described as a dual-specificity protein phosphatase (2). The main substrates of PTEN are inositol phospholipids generated by the activation of the phosphoinositide 3-kinase (PI3K) (3). PTEN is a major negative regulator of the PI3K/Akt signaling pathway (1,4,5). PTEN possesses a carboxy-terminal, noncatalytic regulatory domain with three phosphorylation sites (Ser380, Thr382, and Thr383) that regulate PTEN stability and may affect its biological activity (6,7). PTEN regulates p53 protein levels and activity (8) and is involved in G protein-coupled signaling during chemotaxis (9,10).

To place an order, please [Click HERE](#).