

CDC25C

Reactivity: Human Mouse

Tested applications: WB IF

Recommended Dilution: WB 1:500 - 1:2000 IF 1:50 - 1:200

Calculated MW: 53kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human CDC25C

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CDC25; PPP1R60;

Catalog #: A11328

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 995

Isotype: IgG

Swiss Prot: P30307

Purity: Affinity purification

For research use only.

Background:

Cdc25 is a protein phosphatase responsible for dephosphorylating and activating cdc2, a crucial step in regulating the entry of all eukaryotic cells into mitosis (1). cdc25C is constitutively phosphorylated at Ser216 throughout interphase by c-TAK1, while phosphorylation at this site is DNA damage-dependent at the G2/M checkpoint (2). When phosphorylated at Ser216, cdc25C binds to members of the 14-3-3 family of proteins, sequestering cdc25C in the cytoplasm and thereby preventing premature mitosis (3). The checkpoint kinases Chk1 and Chk2 phosphorylate cdc25C at Ser216 in response to DNA damage (4,5).

To place an order, please [Click HERE](#).