

TIMP2

Reactivity: Human Mouse Rat

Tested applications: WB IHC IF

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200

Calculated MW: 24kDa

Observed MW: Refer to Figures

Immunogen:

Recombinant protein of human TIMP2

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

CSC-21K; TIMP2;

Catalog #: A1558

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 7077

Isotype: IgG

Swiss Prot: P16035

Purity: Affinity purification

For research use only.

Background:

TIMPs are members of the family of tissue inhibitor of matrix metalloproteinases (MMPs) that includes TIMP1, TIMP2, TIMP3, and TIMP4. The main function of TIMPs is their inhibitory effect on MMPs. TIMPs irreversibly inactivate MMPs by direct binding to their catalytic zinc cofactor and resultant inhibition of proteinase function (1,2). In addition to MMP inhibition, TIMPs have also been shown to interact with various membrane receptors on the cell surface. Some of these interactions include: TIMP1 with CD63, TIMP2 with 31 integrin, and TIMP3 with VEGFR2, all of which result in distinct cellular effects (3). TIMPs are involved in a wide variety of biological functions, such as tumor angiogenesis and progression (4,5), wound healing, and vascular remodeling (6,7). Mutations in TIMP3 are associated with Sorsby's fundus dystrophy (8,9).

To place an order, please [Click HERE](#).