

CCL4

Reactivity: Human

Tested applications: WB IHC

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:50 - 1:200

Calculated MW: 10kDa

Observed MW: Refer to Figures

Immunogen:

A synthetic peptide of human CCL4

Storage Buffer:

Store at -20. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

ACT2; G-26; HC21; LAG1; LAG-1; MIP1B; SCYA2; SCYA4; MIP1B1; AT744.1; MIP-1-beta;

Catalog #: A1671

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 6351

Isotype: IgG

Swiss Prot: P13236

Purity: Affinity purification

For research use only.

Background:

Chemokines are members of a superfamily of small inducible, secreted, pro-inflammatory cytokines. Members of the chemokine family exhibit 20 to 50% homology in their predicted amino acid sequences and are divided into four subfamilies. In C-C (or b) subfamily, the first two cysteines are adjacent. C-C chemokines are chemoattractants and activators for monocytes and T cells. C-C subfamily members include macrophage inflammatory protein (MIP)-1 α , MIP-1, MIP-2, MIP-3 α , MIP-3, MIP-4, HCC-1, MIP-5 (or HCC-2), RANTES, MCP-1/2/3 (and the murine homologs JE and MARC), I-309, murine C10 and TCA3. Research has shown that MIP-1 is more selective than MIP-1 α , primarily attracting CD4⁺ T lymphocytes, with a preference for T cells of the naive phenotype. MIP-1 α is a more potent lymphocyte chemoattractant than MIP-1 and exhibits a broader range of chemoattractant specificities. It has been suggested that CD8⁺ T lymphocytes are involved in the control of HIV infection in vivo by the release of HIV-suppressive factors (HIV-SF). MIP-1 α has been identified as one of the major HIV-SFs produced by CD8⁺ T cells, along with MIP-1 and RANTES. Recombinant human MIP-1 α acts as an inhibitor of different strains of HIV-1, HIV-2 and SIV infection in a dose-dependent manner.

To place an order, please [Click HERE](#).