

PSMD5

Reactivity: Human Mouse Rat

Tested applications: WB IHC IP

Recommended Dilution: WB 1:500 - 1:2000 IHC 1:20 - 1:200 IP 1:20 - 1:50

Calculated MW: 56kDa

Observed MW: Refer to figures

Immunogen:

A synthetic Peptide of human PSMD5

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonym:

S5B;

Catalog #: A4056

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 5711

Isotype: IgG

Swiss Prot: Q16401

Purity: Affinity purification

For research use only.

Background:

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits.

Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. This gene encodes a non-ATPase subunit of the 19S regulator base that functions as a chaperone protein during 26S proteasome assembly.

To place an order, please [Click HERE](#).