

HLA-DRB5

Reactivity: Human Mouse Rat

Tested applications: WB IHC

Recommended Dilution: WB 1:200 - 1:2000 IHC 1:20 - 1:200

Calculated MW: 30kDa

Observed MW: Refer to figures

Immunogen:

A synthetic Peptide of human HLA-DRB5

Storage Buffer:

Store at 4. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Catalog #: A9475

Antibody Type:

Polyclonal Antibody

Species: Rabbit

Gene ID: 3127

Isotype: IgG

Swiss Prot: Q30154

Purity: Affinity purification

For research use only.

Background:

HLA-DRB5 belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. DRB1 is expressed at a level five times higher than its paralogues DRB3, DRB4 and DRB5. The presence of DRB5 is linked with allelic variants of DRB1, otherwise it is omitted. There are 4 related pseudogenes: DRB2, DRB6, DRB7, DRB8 and DRB9. [provided by RefSeq, Jul 2008]

To place an order, please [Click HERE](#).