

IL 20 Human

Description: Interleukin-20 Human Recombinant produced in E.Coli is a single, non-glycosylated homodimeric polypeptide chain containing 2 x 153 amino acids and having a total molecular mass of 35,212 Dalton. The IL-20 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-354

For research use only.

Synonyms: IL10D, ZCYTO10, IL-20, MGC96907, Four alpha helix cytokine Zcyto10.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Leu-Lys-Thr-Leu.

Purity: Greater than 95.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Lyophilized carrier free.

Stability:

Lyophilized Interleukin-20 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL20 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin -20 in sterile 18M-cm H2O at a concentration of 0.5mg/ml or less.

Introduction:

IL20 is a cytokine structurally related to interleukin 10 (IL10). IL20 cytokine has been shown to transduce its signal through signal transducer and activator of transcription 3 (STAT3) in keratinocytes. A specific receptor for this cytokine is found to be expressed in skin and upregulated dramatically in psoriatic skin, suggesting a role for this protein in epidermal function and psoriasis.

Biological Activity:

The Biological Activity was determined by measuring the dose dependent activation of STAT3 in human epidermal keratinocytes (HEKa cells). Significant STAT3 activation is observed with 100 ng/mL of recombinant human IL-20 corresponding to a Specific Activity of 10,000IU/mg.

To place an order, please [Click HERE](#).