

IL 10 Porcine

Description: Interleukin-10 Porcine Recombinant produced in E.coli is a single, glycosylated polypeptide chain containing 160 amino acids and having a molecular mass of 18458 Dalton. The IL-10 is purified by proprietary chromatographic techniques.

Catalog #: CYPs-411

Synonyms: B-TCGF, CSIF, TGIF, IL-10, IL10A, MGC126450, MGC126451, Cytokine synthesis inhibitory factor.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: The sequence of the first five N-terminal amino acids was determined and was found to be Met-Ser-Ile-Lys-Ser.

Purity: Greater than 95.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The protein was lyophilized with no additives.

Stability:

Lyophilized Interleukin-10 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL10 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin 10 in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

IL10 is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immunoregulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II Ags, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immunoregulator in the intestinal tract.

Biological Activity:

The ED50 as determined by the dose-dependant stimulation of MC/9 cell which was found to be < 15ng/ml corresponding to a Specific Activity of 66,667IU/mg.

To place an order, please [Click HERE](#).