

Leptin Rat, PEG

Description: Mono-Pegylated Leptin Rat Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 147 amino acids and an additional Ala at N-terminus having a molecular mass of 35.6 kDa (with 20 kDa PEG) as determined by mass spectrometry. However due to enlarged hydrodynamic volume it runs on the SDS-PAGE as 48 kDa protein and in gel-filtration on Superdex 200 as over 100 kDa protein. Its half-life in circulation after SC injection was over 20 hours. Rat Leptin was purified by proprietary chromatographic techniques according to Salomon et al (2006) Protein Expression and Purification 47, 128136 and then pegylated.

Catalog #:CYP5-599

For research use only.

Synonyms: OB Protein, Obesity Protein, OBS, Obesity factor.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Purity: Greater than 99.0% as determined by: (a) Analysis by Gel-Filtration. (b) Analysis by SDS-PAGE.

Formulation:

The protein was lyophilized from a concentrated (0.65mg/ml) solution with 0.003mM NaHCO₃.

Stability:

Lyophilized Rat Leptin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Rat Leptin should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized pegylated Rat Leptin in sterile 0.4% NaHCO₃ adjusted to pH-8.5, not less than 100

Introduction:

A 16-kDa peptide hormone secreted from white adipocytes and implicated in the regulation of food intake and energy balance. Leptin provides the key afferent signal from fat cells in the feedback system that controls body fat stores.

Biological Activity:

Pegylated Rat Leptin is capable of stimulating proliferation of BAF/3 cells stably transfected with the long form of human leptin receptor. Its in vitro activity is only slightly lower than the non-pegylated antagonist but in vivo it has profound weight reducing effect (as compared to the non-pegylated leptin), resulting mainly from reduced food intake.

To place an order, please [Click HERE](#).