

MIA2 Human

Description: MIA2 Human Recombinant produced in E.Coli is a single, non-glycosylated, polypeptide chain consisting of 101 amino having a total molecular mass of 11.5 kDa. The MIA2 is purified by proprietary chromatographic techniques.

Catalog #: CYP5-645

For research use only.

Synonyms: MIA-2, MIA2, Melanoma Inhibitory Activity 2, FLJ22404.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MLESTKLLAD LKKCGDLECE ALINRVSAMR DYRGPDCRYL
NFTKGEEISV YVKLAGERED LWAGSKGKEF GYFPRDAVQI EEVFISEEQ MSTKESDFLC L.

Purity: Greater than 97.0% as determined by (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

The MIA2 protein was lyophilized from a concentrated (1mg/ml) solution containing 20mM Phosphate buffer pH=7.4 and 150mM NaCl.

Stability:

Lyophilized MIA2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution MIA2 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized MIA2 in sterile 18M-cm H₂O not less than 100µg/ml, which can then be further diluted to other aqueous solutions.

Introduction:

MIA2 performs as a tumour suppressor in hepatocellular carcinoma. MIA2 is a novel acute phase protein which its expression is found in the liver and reacts to liver damage in chronic liver diseases and is considerably higher in severe fibrosis patients. MIA2 is induced by IL6, TGF-β & and conditioned medium from activated hepatic stellate cells. MIA2 is mainly expressed in hepatocytes.

To place an order, please [Click HERE](#).