

IL 17B Human

Description: Interleukin-17B Human Recombinant produced in E.Coli is a homodimeric, non-disulfide-linked polypeptide chain containing a total of 322 amino acids (2 chains of 161aa) and having a molecular mass of 36.5kDa. The IL-17B is purified by proprietary chromatographic techniques.

Synonyms: Interleukin-17B, IL-17B, Cytokine Zcyto7, Interleukin-20, Neuronal interleukin-17-related factor, IL20, NIRF, ZCYTO7.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Amino Acid Sequence: MQPRSPKSKR KGQGRPGPLA PGPHQVPLDL VSRMKPYARM
EEYERNIEEM VAQLRNSSSEL AQRKCEVNLQ LWMSNKRSLP PWGYSINHDP SRIPVDLPEA
RCLCLGCVNP FTMQEDRSMV SVPVFSQVPV RRRLCPPPPR TGPCRQRAVM ETIAVGCTCI F.

Purity: Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Formulation:

Interleukin-17B was Lyophilized from a 0.2

Stability:

Lyophilized Interleukin 17B although stable at room temperature for 3 weeks, should be stored desiccated below -18C. Upon reconstitution IL17B should be stored at 4C between 2-7 days and for future use below -18C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Solubility:

It is recommended to reconstitute the lyophilized Interleukin 17B in sterile 18M-cm H2O not less than 100

Introduction:

IL-17 family members are glycoproteins secreted as dimers which induce local cytokine production and recruit granulocytes to sites of inflammation. The IL-17 family is comprised of at least six pro-inflammatory cytokines that share a conserved cysteine-knot structure but diverge at the N-terminus. IL-17 is induced by IL-15 and IL-23, mostly in activated CD4+ T cells distinct from Th1 or Th2 cells. IL-17B binds the IL-17B receptor, but not the IL-17 receptor; it is most homologous with IL-17D, which is expressed by resting CD4+ T cells and CD19+ B cells. IL17B Diseases associated with IL17B include spondyloarthritis, and neuronitis, and among its related super-pathways are Mucin expression in CF via IL-6, IL-17 signaling pathways and STAT3 Pathway.

Biological Activity:

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

The ED50 as determined by inducing IL-8 secretion of human HepG2 cells is less than 1.0



To place an order, please [Click HERE](#).

Catalog #:CYP5-778

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)

Copyrights © 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)