

UCHL3 Human

Description: UCHL3 produced in E.Coli is a single, non-glycosylated polypeptide chain containing 250 amino acids (1-230a.a.) and having a molecular mass of 28.3kDa. UCHL3 is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: ENPS-064

Synonyms: Ubiquitin Carboxyl-Terminal Esterase L3 (ubiquitin thiolesterase), UCH-L3, Ubiquitin Carboxyl-Terminal Hydrolase Isozyme L3, EC 3.4.19.12.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MEGQRWLPLE ANPEVTNQFL
KQLGLHPNWQ FVDVYGMDE LLSMVPRPVC AVLLLPITE KYEVFRTEEE EKIKSQGQDV
TSSVYFMKQT ISNACGTIGL IHAIANNDK MHFESGSTLK KFLEESVSMS PEERARYLEN
YDAIRVTHET SAHEGQTEAP SIDEKVDLHF IALVHVDGHL YELDGRKPPF INHGETSDET
LLEDAIEVCK KF

Purity: Greater than 95% as determined by SDS-PAGE.

Formulation:

The UCHL3 protein solution (1mg/1ml) is formulated in 20mM Tris-HCl buffer (pH8.0) 1mM DTT and 10% glycerol.

Usage:

NeoBiolabs products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Ubiquitin carboxyl-terminal hydrolase isozyme L3 belongs to a gene family whose products hydrolyze small C-terminal adducts of ubiquitin to produce the ubiquitin monomer. UCHL3 takes part in the regulation of neuronal development and spermatogenesis and is associated to neurodegenerative diseases. UCHL3 has a 54% homology to UCHL1.

Biological Activity:

Specific activity: >3,000 pmole/min/μg. Measured by the hydrolysis of Ubiquitin-AMC at pH 8.0, at 37C.

Storage:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Please avoid freeze thaw cycles.

To place an order, please [Click HERE](#).