

MAT2A Human

Description: MAT2A Human Recombinant fused with His tag (20 a.a.) at C-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 415 amino acids and having a molecular mass of 45.8 kDa. The MAT2A is purified by proprietary chromatographic techniques.

Catalog #: ENPS-327

Synonyms: MATA2, MATII, SAMS2, MAT-2A, S-adenosylmethionine synthetase isoform type-2, AdoMet synthetase 2, Methionine adenosyltransferase 2, Methionine adenosyltransferase II, MAT2A, AMS2.

For research use only.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHH SSGLVPRGSH MNGQLNGFHE AFIEEGTFLF
TSESVGEGHP DKICDQISDA VLDAHLQQDP DAKVACETVA KTG MILLAGE ITSRAAVDYQ
KVVREAVKHI GYDDSSKGFD YKTCNVLVAL EQQSPDIAQG VHLDRNEEDI GAGDQGLMFG
YATDETEECM PLTIVLAHKL NAKLAELRRN GTLPWLRPDS KTQVTQYMQ DRGAVLPIRV
HTIVISVQHD EE

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

The MAT2A solution (1mg/ml) contains 20mM Tris pH-8 & 10% glycerol.

Stability:

MAT2A although stable 4°C for 4 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

MAT2A is an important enzyme in cellular metabolism and catalyzes the formation of S-adenosylmethionine (SAME) from L-methionine and ATP. MAT2A is expressed in extrahepatic tissues. In liver, MAT2A expression associates with growth, dedifferentiation, and cancer. NF-kappa B and AP-1 are necessary for basal MAT2A expression in HepG2 cells and mediate the increase in MAT2A expression in response to TNF-alpha. Up-regulation of MAT2A provides growth improvement and s-adenosylmethionine and methylthioadenosine thus can block mitogenic signaling in colon cancer cells. Lower expression of both MAT2A and MAT2beta and interfere with leptin signaling in liver cancer cells.

To place an order, please [Click HERE](#).