

HIV-1 p24 Core, Sf9

Description: HIV-1 p24 Core Recombinant- derived from HIV-1 gene fragment is a 35 kDa gag polyprotein that includes all of the p24 plus additional amino acids of the C-terminus of p17 and the N-terminus of p15. The HIV-1 p24 core is glycosylated with N-linked sugars and produced using baculovirus vectors in insect cells. Purified under conditions that maintain the tertiary structure of the p24 molecule.

Source: Baculovirus Insect Cells.

Physical Appearance: Sterile filtered colorless clear solution.

Purity: Greater than 90.0% as determined by HPLC analysis & SDS-PAGE.

Specificity:

Immunoreactive with sera from HIV infected individuals.

Formulation:

The protein solution contains 10mM Tris pH-8.0, 300mM NaCl and 0.01% Tween-20.

Stability:

Recombinant HIV -1 p24 core sf9 although stable at 4°C for 3 weeks, should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Applications:

HIV-1 p24 Core antigen is suitable for ELISA and Western blots, excellent antigen for early detection of HIV seroconvertors with minimal specificity problems.

Introduction:

Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus *Lentivirus*, part of the family of *Retroviridae*. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the

www.neobiolab.com

info@neobiolab.com

888.754.5670, +1 617.500.7103 United States

0800.088.5164, +44 020.8123.1558 United Kingdom

infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.



To place an order, please [Click HERE](#).

Catalog #:HIPS-160

For research use only.

© 2015 NeoScientific



neobiolab | [Term](#) | [Quote/Order](#) | [Services](#) | [Products](#) | [Support](#) | [Corporate](#) | [Contact Us](#)

Copyrights © 2013 NEO Group 245 First Street, 18th Floor, Cambridge MA 02142 888.754.5670 (toll-free)



Request a
CATALOG
[CLICK HERE](#)