

HIV-1 gp120 CM

Description: HIV-1 gp120 CM Recombinant- is the external envelope protein, full-length 120 kDa, derived from the env. gene of HIV-1 and glycosylated with N-linked sugars and produced using baculovirus vectors in insect cells. The gp120CM stands for Chaing Mai from Clade E and is M-tropic meaning it binds to CCR5. Purified under conditions that maintain the tertiary structure of the molecule.

Source: Baculovirus, Insect Cells.

Physical Appearance: Sterile filtered colorless clear solution.

Purity: Greater than 90.0% as determined by HPLC analysis and SDS-PAGE.

Formulation:

(311

Stability:

Recombinant HIV -1 gp120 CM although stable at 4°C for 3 weeks, should be stored below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Human immunodeficiency virus (HIV) is a retrovirus that can lead to a condition in which the immune system begins to fail, leading to opportunistic infections. HIV primarily infects vital cells in the human immune system such as helper T cells (specifically CD4+ T cells), macrophages and dendritic cells. HIV infection leads to low levels of CD4+ T cells through three main mechanisms: firstly, direct viral killing of infected cells; secondly, increased rates of apoptosis in infected cells; and thirdly, killing of infected CD4+ T cells by CD8 cytotoxic lymphocytes that recognize infected cells. When CD4+ T cell numbers decline below a critical level, cell-mediated immunity is lost, and the body becomes progressively more susceptible to opportunistic infections. HIV was classified as a member of the genus *Lentivirus*, part of the family of *Retroviridae*. Lentiviruses have many common morphologies and biological properties. Many species are infected by lentiviruses, which are characteristically responsible for long-duration illnesses with a long incubation period. Lentiviruses are transmitted as single-stranded, positive-sense, enveloped RNA viruses. Upon entry of the target cell, the viral RNA genome is converted to double-stranded DNA by a virally encoded reverse transcriptase that is present in the virus particle. This viral DNA is then integrated into the cellular DNA by a virally encoded integrase so that the genome can be transcribed. Once the virus has infected the cell, two pathways are possible: either the virus becomes latent and the infected cell continues to function, or the virus becomes active and replicates, and a large number of virus particles are liberated that can then infect other cells.

References:


Title: Slit2/Robo4 Signaling Modulates HIV-1 gp120-Induced Lymphatic

Hyperpermeability. Publication: Zhang X, Yu J, Kuzontkoski PM, Zhu W, Li DY, et al. (2012)

Slit2/Robo4 Signaling Modulates HIV-1 gp120-Induced Lymphatic Hyperpermeability. PLoS

www.neobiolab.com
info@neobiolab.com
888.754.5670, +1 617.500.7103 United States
0800.088.5164, +44 020.8123.1558 United Kingdom
Pathog 8(1): e1002461.

doi:10.1371/journal.ppat.1002461Link:http://www.plospathogens.org/article/info%3Adoi%2F10.1371%2Fjournal.ppat.1002461



Catalog #:HIPS-161

To place an order, please Click [HERE](#).

For research use only.

© 2015 NeoScientific

