

PSMB8 Human

Description: PSMB8 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 229 amino acids (73-276 a.a.) and having a molecular mass of 25.4kDa. PSMB8 is fused to a 25 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: enPS-677

For research use only.

Synonyms: D6S216, D6S216E, LMP7, MGC1491, PSMB5i, Low molecular mass protein 7 Macropain subunit C13, Multicatalytic endopeptidase complex subunit C13, Proteasome component C13, Proteasome subunit beta-5i, Really interesting new gene 10 protein.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered clear solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSMTTTLA FKFQHGVIAA
VDSRASAGSY ISALRVNKVI EINPYLLGTM SGCAADCQYW ERLLAKECRL YYLRNGERIS
VSAASKLLSN MMCQYRGMGL SMGSMICGWD KKGPGLYYVD EHGTRLSGNM FSTGSGNTYA
YGVMDSGYRP NLSPEEAYDL GRRAIAYATH RDSYSGGVVN MYHMKEDGWV KVESTDVSDL
LHQYREANQ.

Purity: Greater than 85.0% as determined by SDS-PAGE.

Formulation:

PSMB8 protein solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.4M Urea and 10% glycerol.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Proteasome Subunit Beta Type 8 (PSMB8) is a member of the peptidase T1B family. The proteasome is a multicatalytic proteinase complex with well-organized ring-shaped 20S core structure. Proteasomes are spread through eukaryotic cells at a high concentration. PSMB8 is mainly expressed in immature dendritic cells. vital function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMB8 is positioned in the class II region of the MHC (main histocompatibility complex). PSMB8 plays as a significant factor of interferon gamma-induced sensitivity and in apoptosis by degradation of the apoptotic inhibitor MCL1.

To place an order, please [Click HERE](#).